



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A fossil Petal and a fossil Fruit from the Cretaceous (Dakota Group) of Kansas

BY ARTHUR HOLLICK

Included in a collection of fossil plant remains from the Cretaceous (Dakota Group) of Kansas, recently obtained by the New York Botanical Garden from Mr. Charles H. Sternberg, of Lawrence, Kans., are two exceedingly interesting specimens — one representing a large petal, the other a fleshy fruit.

Petals, as fossils, are exceedingly rare, and I am not acquainted with any published figure of anything of the kind which can compare with ours, in regard to either size or satisfactory condition of preservation. Unfortunately, a portion of the upper part, including the apex, is gone, but it is sufficiently perfect to indicate what was its original shape, and the principal characters of the nervation are plainly discernible.

Careful examination and comparison has shown that, in all essentials, it agrees with the petals of some of our large-flowered magnolias, such as *M. foetida* Sarg. and *M. macrophylla* Michx., and as some ten species of *Magnolia*, founded upon more or less well-defined leaves, have been described from the Dakota group, I have decided to refer it to that genus.

The probabilities, of course, are that it belongs to a species to which one of the fossil leaves belongs, but as it is impossible definitely to connect the petal with any one of these a distinctive name is necessary, which should indicate that the fossil was a petal and not a leaf.

***Magnolia palaeopetala* sp. nov.**

Petal apparently ovate-spatulate in outline, about 15 cm. long by 10 cm. broad, convex, laterally constricted and incurved at the base; margin entire, wavy or flexuous; nervation flabellate, dictyodrome, simple and well defined below, forking and thinner above, anastomosing, the areolae and reticulations becoming successively smaller and the nervilles finer, towards the margin. (*Fig. A.*)

Locality : Ellsworth Co., Kans.

It is apparent that this petal must have possessed a texture more or less coriaceous, as otherwise its preservation in such a relatively coarse medium as the Dakota sandstone would have been impossible, and this supposition is supported by the fact that the convexity of the surface, the incurving of the base and margin and the constriction of the former, all appear to have been characters of the living flower and not to be due to subsequent distortion by fossilization.



FIG. A. *Magnolia palaeopetalu* Hollick.

The fruit is plainly that of a fig, and although some twenty-three species of *Ficus* have been described from the Dakota group they were based upon leaf-impressions alone. Only two specimens of fruit were known and these were too ill-defined to admit of specific description. ("Fruits of *Ficus*," Lesq. Fl. Dak. Group, 85. *pl.* 10. *f.* 7, 8. See our Fig. B, 2, 3.) They are incidentally mentioned however as being associated with leaves of *Ficus inaequalis* Lesq.

Specimens considerably more satisfactory were described and figured by Heer in 1874, from the lower Atane beds of Greenland — representing a geological horizon which is practically the equivalent of the Dakota group — under the name *Ficus protogaea* Heer* (Fl. Foss. Arct. 3²: 108. *pl.* 30. *f.* 5-7. See our Fig. B, 4, 5, 6.)




FIG. B. 1. *Ficus neurocarpa* Hollick. 2, 3. "Fruits of *Ficus*" (after Lesqueux). 4-6. *Ficus protogaea* Heer (after Heer).

* The name *Ficus protogaea* was previously used by Ettingshausen, in 1867, in describing a fragmentary fossil leaf from the Cretaceous of Niederschoena in Saxony (Sitzb. Akad. Wiss. Wien, Math.-Nat. Cl. 55: 249. *pl.* 2. *f.* 5.) In any revision of the genus it would therefore be necessary to rename these fruits. In this paper however it is not thought necessary to do anything more than to call attention to the fact.

These five specimens are the only ones which I have found recorded as occurring in the same geological horizon as that to which our specimen belongs, or in any strata which may be regarded as its equivalent, and none of them is equal to ours as a specimen, so far as may be judged from the figures.

***Ficus neurocarpa* sp. nov.**

Fruit broadly obovate-spatulate in outline, 2.8 cm. long by 2.4 cm. maximum width, thick and wrinkled towards the middle, flattened more or less at or near the margin, minutely punctate and delicately nerved; primary nerves convergent but distinct and separated from each other at the base, divergent and becoming thinner and indistinct above, where they anastomose; secondary nerves very fine and apparently irregularly disposed. (*Fig. B, 1.*)

Locality: Ellsworth Co., Kans.

This fossil has very much the appearance of many dried herbarium specimens and it is evident that it must have possessed considerable consistency in order to retain its original shape, as it has done to a certain extent, under the pressure to which it must have been subjected.